CEDRIG CLIMATE, ENVIRONMENT AND DISASTER RISK REDUCTION INTEGRATION GUIDANCE



Horti-Sempre Phase 2, Nacala Corridor in Northern Mozambique

Michael Fink, Fabian Mauchle June 2018r.







СЕДРИГ облегченный

Информация общего характера

Contributors Michael Fink, Swisscontact

Fabian Mauchle, SDC, Швейцария

FRANCO SCOTTI, SWISSCONTACT, Бразилия

Tobias Sommer, SDC, Швейцария

Roberto Méndez, COSUDE-Ayuda Humanitaria, Боливия

Общая цель The overall objective of Horti-Sempre is to increase the annual net income of 25,000

smallholders by 30% against baseline by supporting the growth of the horticultural sector in Northern Mozambique in view of its proven importance as income creator.

Страна Мозамбик

Бюджет 6'500'000 CHF

Продолжительность 01/2017 - 12/2020 (48 months)

Аннотация

Описание

The overall objective of the Horti-Sempre Phase 2 Project is to increase smallholder's annual net income by 30% against baseline by supporting the growth of the horticultural sector in Northern Mozambique in view of its proven importance as income creator. To fulfil its mission and reach the overall objective, Swisscontact proposes for Horti-sempre Phase 2 a logic of intervention based on three main Outcomes that unfold around three main project components namely (1) inputs and practices, (2) irrigation and (3) sector competitiveness. OUTCOME No 1: Productivity of horticultural smallholders in the Nacala Corridor in Northern Mozambique increased OUTCOME No 2: Horticultural smallholders in the Nacala Corridor in Northern Mozambique increased their area under irrigation OUTCOME No 3: Market responsiveness and competitiveness of the horti-cultural sector in Northern Mozambique is increased The three components will be complemented with two transversal topics: Women's Economic Empowerment (WEE) throughout the different interventions and through special women targeted interventions and Access to existing funding options. Based on experience from Phase 1, Swisscontact beliefs that Horti-Sempre Phase 2 has the potential to reach 10'000 semi-commercial and 15'000 subsistence male and female smallholders in Northern Mozambique increasing their income by up to 30%.

Сектора, требующие оперативного вмешательства

Сельское хозяйство Развитие сельских районов Продовольственная безопасность Управление водными ресурсами

Документы

MER_Climate Change Profile (pdf, 1.2 Mb)

FANRPAN_Fact Sheet Moz (pdf, 219.89 Kb)

WORLD BANK_Climate Change Profile Moz (pdf, 2.61 Mb)

Presentation_Climate Data_Moz (pdf, 1.01 Mb)

Изображения



Training on basic irrigation solutions

Training on basic irrigation solutions



Training on agricultural practices



Construction of underground dam



Protected horticulture cultivation



Underground dam filled with water

Underground dam filled with water



Basic irrigation solution in use (hip-pump)

Факторы риска

Угрозы, возникающие в результате ухудшения состояния окружающей среды

Деградация (земель, почвы, экосистем, биоразнообразия) Название угрозы Подверженность Да

Soil degradation is frequently the result of poor conservation practices (no soil Комментарии coverage, deep tillage, poor biodiversity) aggravated by heavy rains. Consequently,

more inputs are needed resulting in a vicious circle of degradation.

Последствие Key consequences are lower yields due to degraded soil and higher need of farmers to use inputs (fertilizers)

> Степень вероятности Масштаб Уровень риска Средний ущерб Возможно Средний уровень риска

Загрязнение вод (поверхностных и подземных) Название угрозы

Подверженность Да

Water pollution mainly in peri-urban areas due to urban water and soil Комментарии

contamination (e.g. from factories, waste, etc.).

Key consequences are the loss in product quality, as well as potential health risks for Последствие consumers. Assessment of hazard is difficult because of limited data availability on water and soil quality.

> Степень вероятности Масштаб Уровень риска Небольшой ущерб Возможно Низкий уровень риска

Вредители и эпидемии Название угрозы

Подверженность

Да

Комментарии

Pests and epidemics occur because of poor crop rotation and lack of knowledge or availability of properly formulated defensives (pesticides, insecticides, fungicides). Pests and epidemics are ocurring more frequently during the hot and rainy season compared to the cold and dry season.

Последствие

Key consequences are crop losses (sometimes failure) and that farmers avoid production in warmer and wetter months of the year

Степень вероятности Масштаб Уровень риска СЕДРИГ облегченный

Возможно Средний ущерб Средний уровень риска

Природные угрозы (гидрометеорологические и геологические)

| | геологические) | | | |
|-------------------------------|---|--|---|--|
| Название угрозы | Аномально высока | я температура | | |
| Подверженность | Да | | | |
| Комментарии | According to the World Bank, the number of hot days per year increased by 25 in last 40 years, and much of this has occurred during the southern hemisphere autumn. This corresponds to the first harvest cycle of many major grains across country, with significant implications for agricultural pests and yields. | | e southern hemisphere many major grains across the | |
| Последствие | Key consequences include or crop losses (lower yield | uences include a shortening of the growing season, crop failure (no yield) es (lower yields) | | |
| | Степень вероятности Весьма вероятно | Масштаб Средний ущерб | Уровень риска Высокий уровень риска | |
| | | | | |
| Название угрозы | Засуха | | | |
| Название угрозы | Засуха Не уверен | | | |
| | He уверен Since the 1960s, mean rain month (3.1%) per decade. In variable conditions in the c | ncreased rainfall over the need tentral regions, and persisted south. In Northern Mozamb | erage of 2.5 millimeters per northern regions, highly ent drought periods coupled pique, seasonal droughts are | |
| Подверженность | He yверен Since the 1960s, mean rains month (3.1%) per decade. It variable conditions in the with episodic floods in the occurring, meaning that rains | ncreased rainfall over the need read regions, and persisted south. In Northern Mozamb | northern regions, highly ent drought periods coupled | |
| Подверженность Комментарии | He ybepeh Since the 1960s, mean rain month (3.1%) per decade. It variable conditions in the owith episodic floods in the occurring, meaning that rain Delayed rains result in los | ncreased rainfall over the need read regions, and persisted south. In Northern Mozamb | northern regions, highly ent drought periods coupled pique, seasonal droughts are | |

Название угрозы Бури, торнадо, ураганы, сильный ветер, пылевые бури Подверженность Комментарии Frequency of storms has increased, but events are seasonally concentrated and farmers normally wait with sowing until the risk has decreased. Воследствие Destruction of basic infrastructure and crops in early stage of growth

Степень вероятности Маловероятно

Возможно

Последствие

Масштаб Средний ущерб Уровень риска Низкий уровень риска

Средний уровень

риска

Название угрозыНаводнения, внезапные паводкиПодверженностьДаКомментарииThe proportion of days with heavy rainfall events has increased by 2.6% per decade according to the World Bank. The number of days with heavy rainfall currently amounts to ~25 per year. However, events are seasonally concentrated and farmers normally wait with sowing until this risk is lower.ПоследствиеDestruction of basic infrastructure and crops in early stage of growth, destruction of trade infrastructure (e.g. bridges and roads)Степень вероятностиМасштабУровень риска

| Название угрозы | Erratic Rains | |
|-----------------|---------------|--|
| Подверженность | Да | |

Kommeнтарии Recently, rainfalls in Northern Mozambique are out of usual patterns which farmers rely on. Tendency towards delayed rainfalls.

Средний ущерб

It is difficult for farmers to predict the start of the rainy season. Due to a delayed start of the rainy season, the growing cycle is postponed into the hot season when it is difficult to produce horticulture. Higher risk of pests due to humidity.

 Степень вероятности
 Масштаб
 Уровень риска

 Весьма вероятно
 Средний ущерб
 Высокий уровень риска

Угрозы, возникающие в результате изменения климата (и изменчивости климата)

| Название угрозы | Общие тенденции к повышению или снижению | |
|-----------------|--|--|

| | среднегодовых температур |
|----------------|---|
| Подверженность | Нет |
| Комментарии | Temperatures have generally increased by 0.6° C over the last fourty years, with particularly pronounced increases observed during the hot season (September - March). This increase has so far not considerably affected horticulture as the vegetables are produced during the drier and cooler winter months (April - August). |

Название угрозы

Изменение частоты и интенсивности экстремальных климатических событий и взаимосвязанных стихийных бедствий (например, периодов экстремально высокой или низкой температуры, наводнений, засух, бурь, ураганов и циклонов)

Подверженность

Нет

Комментарии

Frequency of floods is increasing in the country, but mostly in the South and Centre where Mozambique does not control the dam system on the main river (e.g. Limpopo, Save, etc.). Other events (hurricans, cyclones, etc.) are also concentrated in the South/Centre.

Название угрозы

Смещения времен года

Подверженность

Да

Комментарии

A shift of seasons is observed in Northern Mozambique. Average annual rainfall has remained similiar (or even slightly increased). However, the precipitation patterns have changed. More erratic and locally concentrated rainfall is observed which often results in floods and a shorter growing season.

Последствие

Shorter growing season, longer idle season (hunger period - epoca de fome), unpredictability of sowing time, loss of first seeds (investment), extension of growing season into warmer months, loss of 1 or more production cycles

Степень вероятности Весьма вероятно Масштаб Средний ущерб Уровень риска Высокий уровень риска

Нужна ли углубленная оценка риска?

Да – Углубленная оценка риска нужна

• Воздействие

Оцените воздействие на окружающую среду

| Природоохранная сфера | Экосистемы |
|------------------------------------|---|
| Компонент мероприятия | Underground Dams |
| Воздействие на окружающую среду | Small-scale rainwater retention increasing soil humidity potentially changing the ecosystem; limited additional pollution due to the plastic used to build the dam |
| Природоохранная сфера | Почва |
| Компонент мероприятия | Inputs (Fertilizer & Pesticides) |
| Воздействие на окружающую среду | Use of fertilizer and pesticides by horticulture smallholders is common. However, the used amounts are very limited due to a lack of financial resources. Thus, a small negative impact on the soils can be expected. The Project only gives technical advice following a market-approach and does not directly promote and increased use of fertilizers and pesticides for the horticultural production. |
| | Оцените воздействие на изменение климата |
| Компонент | |

компонент мероприятия

Воздействие на изменение климата

Possibly decreasing emissions of Greenhouse Gases (GHG) due to local horticultural production and shorter transport routes. The international and inter-regional imports might decrease due to a higher availability of locally produced vegetables.

Нужна ли углубленная оценка воздействия?

Нет – Углубленная оценка воздействия не нужна